

Phylogenetic study and identification of *Vibrio splendidus*-related strains based on *gyrB* gene sequences

Frédérique Le Roux^{1,*}, Mélanie Gay¹, Christophe Lambert², Jean Louis Nicolas³,
Manolo Gouy⁴, Franck Berthe¹

¹Laboratoire de Génétique et Pathologie, Institut français de recherche pour l'exploitation de la mer (IFREMER), 17390 La Tremblade, France

²Laboratoire des Sciences de l'Environnement Marin (LEMAR), Université de Bretagne Occidentale (UBO), Place Copernic, technopole Brest-Iroise, 29280 Plouzané, France

³Laboratoire de Physiologie des Invertébrés Marins, IFREMER, 29280 Plouzané, France

⁴Laboratoire de Biométrie et Biologie Evolutive, Unité Mixte de Recherche 5558, Centre National de Recherche Scientifique (CNRS), Université Claude Bernard, Lyon, 43 Boulevard du 11 Novembre 1918, 69622 Villeurbanne cedex, France

ABSTRACT: Different strains related to *Vibrio splendidus* have been associated with infection of aquatic animals. An epidemiological study of *V. splendidus* strains associated with *Crassostrea gigas* mortalities demonstrated genetic diversity within this group and suggested its polyphyletic nature. Recently 4 species, *V. lentus*, *V. chagasii*, *V. pomeroyi* and *V. kanaloae*, phenotypically related to *V. splendidus*, have been described, although biochemical methods do not clearly discriminate species within this group. Here, we propose a polyphasic approach to investigate their taxonomic relationships. Phylogenetic analysis of *V. splendidus*-related strains was carried out using the nucleotide sequences of 16S ribosomal DNA (16S rDNA) and gyrase B subunit (*gyrB*) genes. Species delineation based on 16S rDNA-sequencing is limited because of divergence between cistrons, roughly equivalent to divergence between strains. Despite a high level of sequence similarity, strains were separated into 2 clades. In the phylogenetic tree constructed on the basis of *gyrB* gene sequences, strains were separated into 5 independent clusters containing *V. splendidus*, *V. lentus*, *V. chagasii*-type strains and a putative new genomic species. This phylogenetic grouping was almost congruent with that based on DNA–DNA hybridisation analysis. *V. pomeroyi*, *V. kanaloae* and *V. tasmaniensis*-type strains clustered together in a fifth clade. The *gyrB* gene-sequencing approach is discussed as an alternative for investigating the taxonomy of *Vibrio* species.

KEY WORDS: Phylogenetic · *GyrB* · *Vibrio splendidus* · Polyphyletic · Taxonomy

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Vibrio splendidus (Baumann et al. 1980, Baumann & Schubert 1984) is widely distributed in marine ecosystems (Nealson et al. 1993, Farto et al. 1999). Several strains have been associated with mortalities of larvae or spat of the scallop *Pecten maximus* (Nicolas et al. 1996), the turbot *Scophthalmus maximus* (Gatesoupe et al. 1999) and the oyster *Crassostrea gigas* (Sugumar

et al. 1998, Lacoste et al. 2001, Waechter et al. 2002). An epidemiological study of *V. splendidus* strains associated with *C. gigas* summer mortalities demonstrated the important genetic diversity within this group and suggested its polyphyletic nature (Le Roux et al. 2002). Recently, 4 species (*V. lentus*, *V. chagasii*, *V. pomeroyi* and *V. kanaloae*, phenotypically closely related to *V. splendidus*) have been described (Macian et al. 2001, Thompson et al. 2002, 2003a,b). Given the difficulty in

identifying these species by means of biochemical methods and tests currently available, the development of genomic methods, congruent with a quantitative DNA–DNA hybridisation method, appeared to be necessary in a polyphasic approach towards species delineation.

Although 16S rDNA sequence analysis is a standard method to hierarchically order prokaryotic taxa among the ranks of genera and kingdoms (Garrity & Holt 2001, Ludwig & Klenk 2001, Stackebrandt et al. 2002), results of the 16S rDNA-based analysis often do not correlate with DNA–DNA hybridisation, currently recognised as the master criterion for definition of bacterial species (Fox et al. 1992, Stackebrandt & Goebel 1994). While the ability to translate DNA to protein sequences permits phylogenetic analysis of distantly related strains and more accurate sequence alignment (Gupta 1998), use of protein-coding genes may have several advantages over rDNA. Higher levels of sequence variation allow differentiation of closely related strains (Ochman & Wilson 1987). DNA gyrase subunit, *gyrB*, is a Type II topoisomerase found in bacteria. Since 1995, when universal primers for this gene

became available, several publications have suggested that *gyrB* provides suitable sequence data for bacterial phylogenies, possessing essential attributes such as limited horizontal transmission and presence in all bacterial groups (Yamamoto & Harayama 1995, Watanabe et al. 2001).

In this study, phylogenetic analysis of *Vibrio splendidus*-related strains was carried out by means of biochemical identification, *gyrB* and 16S rDNA-gene sequence analyses and quantitative DNA–DNA hybridisation.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The 19 strains used in this study are presented in Table 1 and were isolated from scallops *Pecten Maximus*, turbot *Scophthalmus maximus* and oysters *Crassostrea gigas* during mortality outbreaks, or purchased from national collections (LMG: Laboratorium voor Mikrobiologie, Gent, Belgium; CIP: Collection de l'Institut Pasteur, Paris, France). Strains were maintained on marine

agar 2216 (Difco) at 4°C or stored frozen in marine broth 2216 (Difco) supplemented with 10% (v/v) glycerol (Sigma) at –80°C. Incubations and sub-cultures were at 20°C on marine agar 2216 (Difco). Strains are maintained in the European Community Reference Laboratory for Molluscs Diseases as part of the collections (Centre de Ressources Biologiques, Laboratoire de Génétique et Pathologie, Institut français de recherche pour l'exploitation de la mer [Ifremer], La Tremblade, France).

Phenotypic characterisation. Phenotypic profiles of strains were determined with classical biochemical tests: Gram-staining (Gram kit; Biomerieux); oxidase (bactident oxidase; Merck); respiratory activity (meat liver medium; Diagnostic Pasteur); commercial kits API 20E and API 50CH (Biomerieux); glucose metabolism (MEVAG, Diagnostic Pasteur); motility; NaCl requirement and tolerance (0, 2, 4, 6, 8, 10% w/v) and temperature tolerance (4, 20, 35 and 40°C) were performed in 1.5% (w/v) peptone broth (Diagnostic Pasteur).

Tests were coded as 1 (positive result) or 0 (negative result) and numerical analysis was performed using

Table 1. *Vibrio* spp. strains used in this study and EMBL database accession no. In cases where 2 different sequences were obtained for a single strain, these are labelled a and b. LMG: Laboratorium voor Mikrobiologie, Gent, Belgium; CIP: Collection de l'Institut Pasteur, Paris, France; *P. m.*: *Pecten maximus*; *S. m.*: *Scophthalmus maximus*; *C. g.*: *Crassostrea gigas*; *N. n.*: *Nodipecten nodosus*; *O. e.*: *Ostrea edulis*

Strain designation	Identification	Source	Accession no.	
			16S	<i>gyrB</i>
LMG 4042 ^T	<i>V. splendidus</i>	Marine fishes	a:AJ515229 b:AJ515230	AJ515292
PMV18	<i>V. splendidus</i>	<i>P. m.</i> larvae	AJ515224	AJ515284
PMV19	<i>V. splendidus</i>	<i>P. m.</i> larvae		AJ577816
A515	<i>V. splendidus</i>	<i>P. m.</i> larvae	a:AJ515222 b:AJ515223	AJ515283
LT06	<i>V. splendidus</i>	<i>P. m.</i> larvae	AJ515221	AJ515282
CIP107166 ^T	<i>V. lentus</i>	<i>O. e.</i>	AJ294421	AJ515293
VS6 turb	<i>V. lentus</i>	<i>S. m.</i> larvae	AJ515228	AJ515288
A053	<i>V. lentus</i>	<i>P. m.</i> larvae	a:AJ515225 b:AJ515226	AJ515286
VC1	<i>V. lentus</i>	<i>P. m.</i> larvae	AJ515227	AJ515287
LMG 21353 ^T	<i>V. chagasii</i>	<i>S. m.</i> larvae	AJ316199	AJ577820
TNEMF6	<i>V. chagasii</i>	<i>C. g.</i> spat 1997	AJ515218	AJ515264
TNNIII7	<i>V. chagasii</i>	<i>C. g.</i> spat 1997	AJ515219	AJ515265
LMG 16745	<i>V. chagasii</i>	Marine fishes		AJ515278
LMG 20537 ^T	<i>V. pomeroiyi</i>	<i>N. n.</i>	AJ491290	AJ577822
LMG 20539 ^T	<i>V. kanaloae</i>	<i>O. e.</i>	AJ316193	AJ577821
LMG 20012 ^T	<i>V. tasmaniensis</i>	Marine fishes	AJ316192	AJ577823
Mel 13	Unknown	<i>C. g.</i> spat 2001		AJ577817
Mel 107	Unknown	<i>C. g.</i> spat 2001		AJ577818
Mel 108	Unknown	<i>C. g.</i> spat 2001		AJ577819

simple matching coefficients (Sneath 1972) and an unweighted pair-group method (Sneath & Sokal 1973).

Extraction of genomic DNA, amplification, cloning and sequencing of DNA. Genomic DNA was prepared according to Sambrook et al. (1989). PCR amplification of 16S rDNA and *gyrB* was achieved with primers, and following methods previously described (Yamamoto & Harayama 1995, Lambert et al. 1998). Amplified products were inserted in the pCR2.1 Vector System (Invitrogen) and used to transform *Escherichia coli* cells. We selected 2 to 4 colonies per strain and gene; minipreps of recombinant plasmids were performed according to standard alkaline-lysis protocols, with an additional phenol/chloroform extraction step and ethanol precipitation (Sambrook et al. 1989). Sequencing was carried out using the Sequiterm Excell II kit (Epicentre) and a Li-cor DNA sequencer (ScienceTec) according to the manufacturer's instruction.

Phylogenetic analyses. 16S rDNA and *gyrB* sequences were aligned and phylogenetic analyses were performed using Seaview and Phylo-win programs (Galtier et al. 1996). Phylogenetic trees were built using the BIONJ method (Gascuel 1997) applied to Kimura's 2-parameter distances. Reliability of topologies was assessed by the bootstrap method with 1000 replicates. When 2 different sequences were obtained for a single strain, they were labelled a and b in the phylogenetic tree.

Quantitative DNA–DNA hybridisations. We used and compared 2 methods for DNA–DNA hybridisation experiments. Labelling of DNA probes with tritium-

labelled nucleotides was performed by the random primer method (Megaprime labelling kit) and hybridisation was carried out at 60°C by the S1-nuclease method (Crosa et al. 1973, Grimont et al. 1980) with adsorption of S1-resistant onto DE81 filters (Whatman). A second hybridisation method was performed by a non-radioactive dot-blot method described previously (Macian et al. 2001, Le Roux et al. 2002). Statistical analyses were performed using Statgraphics software, Version 5.1. The relation between *gyrB* sequencing and DNA–DNA hybridisation (S1-nuclease method) was calculated using a reciprocal model, whereas the comparison of the 2 DNA–DNA hybridisation methods was carried out using a linear model.

RESULTS

Phenotypic characterisation

Phenotypic analysis, based on 80 biochemical tests, confirmed the relatedness of the 19 strains included in this study (70 % similarity). The 19 discriminating tests were: β -galactosidase, arginine dihydrolase, tryptophan desaminase, acetoin production, oxidation/fermentation of saccharose, melibiose, amygdaline, glycerol, galactose, N-acetyl glucosamine, esculine, celibiose, amidon, gluconate, NO₂ production, growth at 4, 35 and 40°C and 6 % NaCl (Table 2). The results are presented as a dendrogram using simple matching-similarity coefficients (Fig. 1).

Table 2. *Vibrio splendidus*. Phenotypic characteristics of the 19 strains examined in this study. Only results of discriminant tests are shown. 1: β -galactosidase; 2: arginine dihydrolase; 3: tryptophan desaminase; 4: acetoin production; oxidation/fermentation of 5: saccharose; 6: melibiose; 7: amygdaline; 8: glycerol; 9: galactose; 10: N-acetyl glucosamine; 11: esculine; 12: celibiose; 13: amidon; 14: gluconate; 15: NO₂ production; growth at 16: 4°C; 17: 35°C; 18: 40°C; 19: 6 % NaCl

Strain designation	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
LMG 4042 ^T	+	-	-	-	-	+	+	+	+	+	+	+	+	-	-	+	+	-	-
PMV18	+	+	-	-	-	-	+	+	+	+	+	+	+	-	+	+	-	-	-
PMV19	+	+	-	-	-	-	+	+	+	+	+	+	+	-	+	+	+	-	-
A515	+	+	+	-	-	+	+	+	+	+	+	+	+	-	+	+	-	-	-
LT06	+	+	+	+	+	-	+	-	+	+	+	+	+	-	+	+	-	-	-
CIP107166 ^T	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
VS6 turb	-	+	-	-	-	-	+	+	+	+	+	+	+	-	+	-	-	-	-
A053	+	+	-	-	-	-	+	-	+	+	-	+	+	-	+	+	-	-	-
VC1	+	+	-	-	-	-	+	-	+	+	+	+	+	-	+	-	-	-	-
LMG 21353 ^T	-	+	+	-	+	-	-	-	-	+	+	+	+	-	-	-	-	-	+
TNEMF6	-	+	+	-	-	-	+	+	+	+	+	+	+	+	+	+	+	-	-
TNNIII7	-	+	+	-	-	-	+	-	+	+	+	+	+	+	+	-	+	-	-
LMG 16745	-	+	-	-	-	-	+	+	+	+	+	+	+	-	+	-	+	-	+
LMG 20537 ^T	+	+	+	-	-	-	+	-	+	+	+	+	+	-	-	+	-	-	+
LMG 20539 ^T	+	+	+	-	+	-	+	-	+	+	+	+	+	-	-	+	+	+	+
LMG 20012 ^T	-	-	-	-	-	-	-	-	-	+	+	+	-	-	+	-	-	-	+
Mel 13	-	+	-	-	-	+	+	+	+	+	+	+	+	-	+	+	-	-	-
Mel 107	-	+	-	-	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
Mel 108	-	+	-	-	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-




Fig. 1. *Vibrio* spp. Cluster-analysis of phenetic data using simple matching-similarity coefficient and unweighted pair-group method (percentage in top abscissa) for 18 strains related to *V. splendidus*

Phylogenetic analysis based on 16S rDNA

The 16S rDNA fragments were amplified and cloned. The sequences used for constructing the phylogenetic tree were 1332 gap-free sites in length. The 4 sequences obtained from each strain were similar, except for A053, A515 and LMG 4042^T, for which polymorphism (2% divergence) was observed. This polymorphism involved an *Hha*1 restriction site, and PCR-RFLP (restriction fragment-length polymorphism) on other clones confirmed this result (data not shown). Despite a high level of sequence similarity (98%), a cluster comprising 3 strains: TNEMF6, TNNIII7 and *Vibrio chagasii*-type Strain LMG 21353^T was supported by a bootstrap value of 96% (Fig. 2). In this phylogenetic tree the *V. lentus* type strain clustered with the *V. splendidus*-type strain, suggesting that 16S rDNA sequence-analysis does not differentiate these 2 species.

Phylogenetic analysis based on *gyrB*

The *gyrB* fragments were amplified and cloned. The sequences used in constructing the phylogenetic tree were 1064 gap-free sites long. Sequences of different cloned fragments obtained from previous strains were similar. The phylogenetic tree based on the *gyrB* nucleotide sequences from *Vibrio splendidus*-related strains confirmed clustering of LMG 21353^T, TNEMF6 and TNNIII7 (Fig. 3). Within this clade, clustering of TNEMF6, TNNIII7 and LMG 16745 was supported by

a bootstrap value of 93%. A second clade, supported by a bootstrap value of 100%, comprised 3 strains isolated from *Crassostrea gigas*: Mel 108, Mel 107 and Mel 13. A third clade, with a bootstrap value of 100%, comprised *V. splendidus*-type strain (LMG 4042^T), LT06, A515, PMV18 and PMV19. A 4th clade comprised *V. lentus*-type strain and A053, VS6turb, and VC1. *V. kanaloae*, *V. pomeroyi* and *V. tasmaniensis*-type strains were clearly separated from this 4th clade, and clustered together in a fifth clade. Similar results were obtained by 3 methods: neighbour-joining, maximum parsimony and maximum-likelihood analysis (data not shown).

Table 3. *Vibrio splendidus*. Intraspecific DNA–DNA homology among strains of 3 clades in *V. splendidus* group determined by (a) S1-nuclease method and (b) the dot-blot method

Test strain	Probe TNEMF6		Probe LMG 4042 ^T		Probe A053	
	a	b	a	b	a	b
LMG 4042 ^T	40	40	100	100	64	56
PMV18		47	75	84		70
A515		37	85	70		54
LT06		44	64	86		57
CIP107166 ^T	39		53		79	
A053		38		53	100	100
VC1		35		50	75	90
TNEMF6	100	100		40		49
TNNIII7	83	92	44	35	39	50
LMG 16745	63	79				




Fig. 2. *Vibrio* spp. Phylogenetic tree of partial 16S rDNA sequences. *V. campbellii* homologue was used as outgroup (EMBL accession no. X56575); 1332 gap-free sites were compared. Horizontal branch lengths are proportional to evolutionary divergence. Bootstrap values (as % of 1000 replicates) appear next to corresponding branch when >50%. In cases where 2 different sequences were obtained for a single strain, these are labelled a and b

DNA–DNA hybridisation

We used 2 methods of DNA–DNA hybridisation, S1-nuclease and dot-blot, to analyse genomic relatedness between isolates and type strains among 3 clades revealed by the phylogenetic tree in Fig. 3.

A first hybridisation was performed with strains LMG 4042^T, CIP 107166^T, TNEMF6, TNNIII7 and LMG 16745 using labelled TNEMF6 DNA as a probe. High rates of hybridisation were obtained with the dot-blot method, with percentages of re-association of 92 and 79% for TNNIII7 and LMG 16745 respectively (Table 3). Using the S1 nuclease method, the rates of hybridisation were 83 and 63% respectively. With both methods the re-association percentage was lower with LMG 4042^T and CIP 107166^T, i.e. 40 to 39% respectively.

A second set of hybridisations was performed with strains LMG 4042^T, PMV18, A515, LT06, CIP 107166^T, TNEMF6 and TNNIII7, using LMG 4042^T as a probe. Again, higher rates of hybridisation were obtained with the dot-blot method with percentages of re-association above 70% for

PMV18, A515 and LT06. Hybridisation with TNEMF6 was lower, i.e. 40%. With the S1 nuclease method, rates of hybridisation for the strains belonging to *Vibrio splendidus* ranged between 64 and 85%, and hybridisation with species from the other clade was between 44 and 53%.

A third set of hybridisations was performed with LMG 4042^T, CIP 107166^T, A053, VC1, TNEMF6 and TNNIII7, using labelled A053 DNA as a probe. The S1-nuclease method showed percentages of re-association above 75% for CIP 107166^T, A053 and VC1. Hybridisation with TNNIII7 was lower, 39%. This result was confirmed by the second method.

Comparison of methods

A reciprocal model was used to compare *gyrB* sequencing and DNA–DNA hybridisation using the S1-nuclease method (Fig. 4a). The adjusted model was: *gyrB* sequence = 109.188 – 917.844/hybridisation. The ANOVA probability value was <0.01, and adjustment between these 2 methods was statistically significant at the




Fig. 3. *Vibrio* spp. Phylogenetic tree of partial *gyrB* sequences. *V. campbellii* homologue was used as outgroup (EMBL accession no. ABO 14950); 1064 gap-free sites were compared. Other features as in Fig. 2




Fig. 4. Comparison of methods. (a) Reciprocal model used to compare *gyrB* sequencing and DNA–DNA hybridisation in *Vibrio splendidus* using S1-nuclease (% hybridisation in abscissa; % sequence similarities in ordinate); (b) linear model comparing S1-nuclease and dot-blot methods for determining DNA–DNA hybridisation (% hybridisation in abscissa and ordinate)

99% confidence level. The coefficient of correlation was 0.86.

A linear model was used to compare the 2 methods of DNA–DNA hybridisation (Fig. 4b). The adjusted equation was $\text{dot-blot} = 17.1881 + 0.851149 \times \text{S1-nuclease}$. The ANOVA probability value was <0.01 , and adjustment between these 2 methods was statistically significant at the 99% confidence level. The coefficient of correlation was 0.84.

DISCUSSION

Different strains related to *Vibrio splendidus* have been increasingly associated with infection of aquatic species (Sugumar et al. 1998, Gatesoupe et al. 1999, Lacoste et al. 2001, Waechter et al. 2002). As aquaculture production is highly vulnerable to the impact of infectious diseases, the prophylactic approach is of central importance. Given this, it is necessary to develop new diagnostic tests to detect specific pathogens in fishes, shellfishes and the environment, and to monitor the spread and evolution of diseases. In this context, definition and identification of bacterial species and strains raise taxonomy to its pivotal role.

Most of the *Vibrio splendidus*-related strains are generally identifiable only by means of phenotypic characterisation. This study has confirmed that the biochemical tests currently available are not sufficiently discriminatory to distinguish between strains of *V. splendidus*, *V. lentus*, *V. chagasii*, *V. pomeroyi* and *V. kanaloae* species. Even though descriptions of new species based on biochemical tests of relatively low numbers of strains may provide a discriminatory basis, these tests prove highly variable when applied to a broader spectrum of strains belonging to this species. This could be due to metabolic versatility of strains

exposed to variations in the marine environment. Bacterial identification procedures should take this into consideration by incorporating DNA-based methods. A polyphasic approach using molecular techniques is now widely recognised and recommended for species definition (Thompson et al. 2001, Stackenbrandt et al. 2002). This should provide new identification tools for use in diagnostic laboratories.

Our preliminary phylogenetic study was based on partial 16S rDNA gene sequences. The results are in accordance with those of previous studies showing that 16S rDNA sequences of *Vibrio splendidus*-related strains are very similar (Macian et al. 2001, Le Roux et al. 2002, Waechter et al. 2002, Thompson et al. 2003a, b). They would confirm the current opinion that 16S rDNA sequences essentially permit phylogenetic analysis of distantly related strains (Gupta 1998) are probably less effective in identifying closely related strains or even species. In our study, only 1 cluster, containing the *V. chagasii*-type strain, was supported by a high bootstrap value. Sequence dimorphism was observed for several strains (Fig. 1), possibly resulting from microheterogeneity between cistrons (Fox et al. 1992), whereby divergence between cistrons is equivalent to divergence between strains. Therefore, these strains cannot be identified from our data. Bearing in mind the above limitations, the use of protein-coding genes as a data source was considered in the following analysis.

Fig. 3 shows the phylogenetic grouping based on *gyrB* sequences. The grouping is supported by strong bootstrap values and almost consistent with the results achieved with the quantitative DNA–DNA hybridisation method (Table 3). The primary indication is that strains phenotypically related to *Vibrio splendidus* can belong to distinct species, which strongly supports the suggested polyphyletic nature of *V. splendidus* and a

diversity of potentially new species related to *V. splendidus* (Macian et al. 2001, Thompson et al. 2001, 2003a,b, Le Roux et al. 2002).

Strains belonging to *Vibrio splendidus*, *V. lentus* and *V. chagasii* species were clearly separated. The delineation of TNEMF6, TNNIII7 and LMG 16745 from *V. chagasii* is suggested by a high bootstrap value in the *gyrB* phylogenetic tree. However, further studies including more strains from these 2 groups and DNA–DNA hybridisation data are necessary to test the validity of our results. Our data also suggest that Mel 108, Mel 107 and Mel 13 belong to a new taxon. Here again, further DNA–DNA hybridisation is needed, and in a further study currently in progress we hope to confirm these results and possibly determine 1 or more new species.

For *Vibrio pomereyi*, *V. kanaloae* and *V. tasmaniensis*, our *gyrB* based analysis appears less discriminative than fluorescent-amplified fragment-length polymorphism (FAFLP) fingerprintings applied to this group (Thompson et al. 2001, 2003a,b). Further work incorporating more strains belonging to these species should be conducted to compare these 2 approaches.

Quantitative DNA–DNA hybridisation is still the recognised reference method for bacterial systematics (Grimont 1984). In our study, we used and compared 2 different methods, namely S1-nuclease and dot-blot. Although the results were well-correlated, the dot-blot method frequently yielded higher rates of re-association than the S1-nuclease method (Table 3). The S1-nuclease method, using enzyme activity to digest single-stranded DNA, is probably more accurate, avoiding counts of partially hybridised probe. Hybridisation rates between 2 strains varied as a function of method (S1-nuclease/dot-blot). This raises the issue of the thresholds currently accepted for species recognition, since values may vary with method employed. For instance, the probe A053 gave a 90% hybridisation rate with VC1 using the dot-blot method, but 75% with the S1-nuclease method. Thus, in view of the fact that dot-blot results probably overestimate the homology and that the S1-nuclease method was not validated by a ΔT_m (evaluation of hybrid stability) calculation (Grimont 1984), VC1 and A053 cannot be considered to comprise a single species. However within the framework of this study, DNA–DNA hybridisations were used as tendency indicators to validate phylogenetic constructions based on *gyrB* gene sequences; e.g. PMV18 displayed 84% hybridisation with *Vibrio splendidus*-type strain compared to 70% for A053 and 47% for TNEMF6, which was in line with the phylogenetic *gyrB* analyses.

A problem associated with quantitative DNA–DNA hybridisation recently arose during sequence analyses of complete genomes. Genomic comparisons revealed

that strains belonging to the same species may contain large amounts (as many as 1.1 Megabasis out of 5) of strain-specific sequences. In the case of *Escherichia coli* strain O157 analysis surprisingly revealed a large number of prophages (Ohnishi et al. 2001). Such numerous strain-specific sequences could strongly alter the hybridisation rates between 2 strains and consequently lead to misidentification of species.

Our results support the proposal of other investigators that *gyrB*-based phylogenetic structure comprises an alternative method to DNA–DNA hybridisation for determining the taxonomic relationships of *Vibrio splendidus*-related strains (Suzuki et al. 1999, Yamamoto et al. 1999). Until the method has been rendered more effective, *gyrB* sequences could be used to cluster strains, and such clusters could then be validated by DNA–DNA hybridisation using S1-nuclease and ΔT_m calculation. Other house-keeping genes, e.g. *rpoD* (Yamamoto & Harayama 1998) and *hsp60* (Kwok et al. 2002), should also be sequenced, incorporating *gyrB* in a multilocus scheme for the future identification and classification of *Vibrio* species. Such sequence information would enable the development of molecular diagnostic tools to detect specific species or strains responsible for mortality events.

Acknowledgements. The authors acknowledge Professor J. Swing for providing LMG type strains, Professor B. Austin and Dr. F. Thompson for critically reading the manuscript, and Dr. P. Soletchnik for statistical analyses. This study was carried out with financial assistance from the European Community Reference Laboratory for Bivalve Mollusc Diseases.

LITERATURE CITED

- Baumann P, Schubert RHW (1984) Family II *Vibrionaceae* Veron 1965, 5245^{AL}. In: Krieg NR, Holt JG (eds) *Bergey's manual systematic of bacteriology*. Williams & Wilkins, Baltimore, p 516–517
- Baumann P, Baumann L, Bang SS, Woolkalis MJ (1980) Reevaluation of the taxonomy of *Vibrio*, *Beneckeia*, and *Photobacterium*: abolition of the genus *Beneckeia*. *Curr Microbiol* 4:127–132
- Crosa JH, Brenner DJ, Falkow S (1973) Use of a single-strand specific nuclease for analysis of bacterial and plasmid deoxyribonucleic acid homo- and heteroduplexes. *J Bacteriol* 115:904–911
- Farto R, Montes M, Perez MJ, Nieto TP, Larsen JL, Pedersen K (1999) Characterization by numerical taxonomy and ribotyping of *Vibrio splendidus* biovar I and *Vibrio scophthalmi* strains associated with turbot cultures. *J Appl Microbiol* 86:796–804
- Fox GE, Wisotzkey JD, Jurtshuk PJ (1992) How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. *Int J Syst Bacteriol* 42:166–170
- Galtier N, Gouy M, Gautier C (1996) SEAVIEW and PHYLO_WIN: 2 graphic tools for sequence alignment and molecular phylogeny. *Comput Appl Biosci* 12:543–548
- Garrity GM, Holt JG (2001) The road map to the manual. In: Boone DR, Castenholz RW, Garrity GM (eds) *Bergey's*

- manual of systematic bacteriology, 2nd edn. Springer-Verlag, New York, p 119–166
- Gascuel O (1997) BIONJ: an improved version of the NJ algorithm based on a simple model of sequence data. *Mol Biol Evol* 14:685–695
- Gatesoupe FJ, Lambert C, Nicolas JL (1999) Pathogenicity of *Vibrio splendidus* strains associated with turbot larvae, *Scophthalmus maximus*. *J Appl Microbiol* 87:757–763
- Grimont PAD (1984) Use of DNA reassociation in bacterial classification. *Can J Microbiol* 34:541–546
- Grimont PAD, Popoff MY, Grimont F, Coynault C, Lemelin M (1980) Reproducibility and correlation study of three deoxyribonucleic acid hybridisation procedure. *Curr Microbiol* 4:325–330
- Gupta RS (1998) Protein phylogenies and signature sequences: a reappraisal of evolutionary relationships among Archaeobacteria, Eubacteria, and Eukaryotes. *Microbiol Mol Biol Rev* 62:1425–1491
- Kwok AY, Wilson JT, Coulthart M, Ng LK, Mutharia L, Chow AW (2002) Phylogenetic study and identification of human pathogenic *Vibrio* species based on partial *hsp60* gene sequences. *Can J Microbiol* 48:903–910
- Lacoste A, Jalabert F, Malham S, Cuffe A, Gélébart F, Cordevant C, Lange M, Poulet SA (2001) A *Vibrio splendidus* strain is associated with summer mortality of juvenile oysters *Crassostrea gigas* in the Bay of Morlaix (North Brittany, France). *Dis Aquat Org* 46:139–145
- Lambert C, Nicolas JL, Cilia V, Corre S (1998) *Vibrio pectenica* sp. nov. a pathogen of scallop (*Pecten maximus*) larvae. *Int J Syst Bacteriol* 48:481–487
- Le Roux F, Gay M, Lambert C, Waechter M, Poubalanne S, Chollet B, Nicolas JL, Berthe F (2002) Comparative analysis of *Vibrio splendidus*-related strains isolated during *Crassostrea gigas* mortality events. *Aquat Living Resour* 15:251–258
- Ludwig W, Klenk HP (2001) Overview: a phylogenetic backbone and taxonomic frame work for prokaryotic systematics. In: Boone DR, Castenholtz RW, Garrity GM (eds) *Bergey's manual of systematic bacteriology*, 2nd edn. Springer-Verlag, New York, p 49–65
- Macian MC, Ludwig W, Aznar R, Grimont PA, Schleifer KH, Garay E, Pujalte MJ (2001) *Vibrio lentus* sp. nov., isolated from Mediterranean oysters. *Int J Syst Evol Microbiol* 51:1449–1456
- Nealson KH, Wimpee B, Wimpee C (1993) Identification of *Vibrio splendidus* as a member of the planktonic luminous bacteria from Persian Gulf and Kuwait region with LuxA probes. *Appl Environ Microbiol* 59:2684–2689
- Nicolas JL, Corre S, Gauthier G, Robert R, Ansquer D (1996) Bacterial problems associated with scallop *Pecten maximus* larval culture. *Dis Aquat Org* 27:67–76
- Ochman H, Wilson AC (1987) Evolution in bacteria: evidence for a universal substitution rate in cellular genomes. *J Mol Evol* 26:74–86
- Ohnishi M, Kurokawa K, Hayashi T (2001) Diversification of *Escherichia coli* genomes: are bacteriophages the major contributors? *Trends Microbiol* 9:481–485
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*, 2nd edn. Cold Spring Harbor, NY
- Sneath PHA (1972) Computer taxonomy. *Methods Microbiol* 7A:29–98
- Sneath PHA, Sokal RR (1973) *Numerical taxonomy: the principles and practice of numerical classification*. WH Freeman, San Francisco, p 188–1308
- Stackebrandt E, Goebel BM (1994) Taxonomic note: a place for DNA–DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* 44:846–849
- Stackebrandt E, Frederiksen W, Garrity GM, Grimont PAD and 9 others (2002) Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *Int J Syst Evol Microbiol* 52:1043–1047
- Sugumar G, Nakai T, Hirata Y, Matsubara D, Muroga K (1998) *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis Aquat Org* 33:111–118
- Suzuki M, Nakagawa Y, Harayama S, Yamamoto S (1999) Phylogenetic analysis of genus *Marinilabilia* and related bacteria based on the amino acid sequences of *gyrB* and amended description of *Marinilabilia salmonicolor* with *Marinilabilia agarovorans* as its subjective synonym. *Int J Syst Bacteriol* 49:1551–1557
- Thompson FL, Hoste B, Vandemeulebroecke K, Swings J (2001) Genomic diversity amongst *Vibrio* isolates from different sources determined by fluorescent amplified fragment length polymorphism. *Syst Appl Microbiol* 24:520–38
- Thompson FL, Thompson CC, Li Y, Gomez-Gil B, Vandenberghe J, Hoste B, Swings J (2003a) *Vibrio kanaloae* sp. nov., *Vibrio pomeroyi* sp. nov. and *Vibrio chagasii* sp. nov., from sea water and marine animals. *Int J Syst Evol Microbiol* 53:753–759
- Thompson FL, Thompson CC, Swings J (2003b) *Vibrio tasmaniensis* sp. nov., isolated from Atlantic salmon (*Salmo salar* L.). *Syst Appl Microbiol* 26:65–69
- Waechter M, Le Roux F, Nicolas JL, Marissal E, Berthe F (2002) Characterisation of *Crassostrea gigas* spat pathogenic bacteria. *C R Acad Sci* 325:231–238
- Watanabe K, Nelson J, Harayama S, Kasai H (2001) ICB database: the *gyrB* database for identification and classification of bacteria. *Nucleic Acids Res* 29:344–145
- Yamamoto S, Harayama S (1995) PCR amplification and direct sequencing of *gyrB* genes with universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. *Appl Environ Microbiol* 61:1104–1109
- Yamamoto S, Harayama S (1998) Phylogenetic relationships of *Pseudomonas putida* strains deduced from the nucleotide sequences of *gyrB*, *rpoD* and 16SrRNA genes. *Int J Syst Bacteriol* 48:813–819
- Yamamoto S, Bouvet PJ, Harayama S (1999) Phylogenetic structures of the genus *Acinetobacter* based on *gyrB* sequences: comparison with the grouping by DNA–hybridisation. *Int J Syst Bacteriol* 49:87–95

Editorial responsibility: Carey Cunningham,
Aberdeen, UK

Submitted: July 28, 2003; Accepted: October 21, 2003
Proofs received from author(s): February 23, 2004